

=> d his full

(FILE 'HOME' ENTERED AT 13:58:35 ON 20 SEP 2001)

FILE 'CAPLUS, EMBASE, MEDLINE, USPATFULL, PCTFULL, EUROPATFULL' ENTERED
AT 13:59:59 ON 20 SEP 2001

L1 577 SEA ABB=ON PLU=ON (TNF OR TUMOR NECRO? FACTOR OR TUMOUR
NECRO? FACTOR) AND (HEARING OR COCHLEAR ADJ CELLS OR AUDITORY)

L2 430 SEA ABB=ON PLU=ON (TNF OR TUMOR NECRO? FACTOR OR TUMOUR
NECRO? FACTOR) AND (HEARING (3A) LOSS OR COCHLEAR (3A)
DEGENRAT? OR AUDITORY)

L3 340 SEA ABB=ON PLU=ON (TNF OR TUMOR NECRO? FACTOR OR TUMOUR
NECRO? FACTOR) (P) (HEARING (3A) LOSS OR COCHLEAR (3A)
DEGENRAT? OR AUDITORY)

L4 404 SEA ABB=ON PLU=ON (TNF OR TUMOR NECRO? FACTOR OR TUMOUR
NECRO? FACTOR) (L) (HEARING (3A) LOSS OR COCHLEAR (3A)
DEGENRAT? OR AUDITORY)

L5 406 SEA ABB=ON PLU=ON (TNF OR TUMOR NECRO? FACTOR OR TUMOUR
NECRO? FACTOR) (L) ((HEARING (3A) LOSS) OR (COCHLEAR (3A)
DEGENERAT?) OR AUDITORY)

L6 3 SEA ABB=ON PLU=ON (TNF OR TUMOR NECRO? FACTOR OR TUMOUR
NECRO? FACTOR) (5A) ((HEARING (3A) LOSS) OR (COCHLEAR (3A)
DEGENERAT?) OR AUDITORY)

L7 3 DUP REM L6 (0 DUPLICATES REMOVED)
D L7 IBIB KWIC
D L7 IBIB KWIC 2-

L8 393 DUP REM L5 (13 DUPLICATES REMOVED)

L9 142 SEA ABB=ON PLU=ON (TNF OR TUMOR NECRO? FACTOR OR TUMOUR
NECRO? FACTOR) (5A) (INHIBIT? OR MODIF? OR ANTAGONIST OR
ANTIBOD?) (P) (HEARING LOSS OR COCHLEAR DEGENERAT?)

L10 142 SEA ABB=ON PLU=ON (TNF OR TUMOR NECRO? FACTOR OR TUMOUR
NECRO? FACTOR) (5A) (INHIBIT? OR MODIF? OR ANTAGONIST OR
ANTIBOD?) (P) (HEARING (3A) LOSS OR COCHLEAR (3A) DEGENERAT?)
D L10 1-5 IBIB KWIC
D L10 130-142 IBIB KWIC

L12 127 SEA ABB=ON PLU=ON L10 (P) (INTERLEUKIN-1 OR IL-1)
D L12 120-127 IBIB KWIC

1Q ANSWER 139 OF 142 PCTFULL COPYRIGHT 2001 MicroPatent
 ACCESSION NUMBER: 1997042966 PCTFULL
 TITLE (ENGLISH): THERAPEUTIC USES OF BPI PROTEIN PRODUCTS FOR HUMAN
 MENINGOCOCCEMIA
 TITLE (FRENCH): APPLICATIONS THERAPEUTIQUES DE PRODUITS PROTEIQUES
 BACTERICIDES/AUGMENTANT LA PERMEABILITE (BPI) DANS LE
 CAS DE
 MENINGOCOCCEMIES CHEZ L'HOMME
 INVENTOR(S): GIROIR, Brett, P.; SCANNON, Patrick, J.
 PATENT ASSIGNEE(S): XOMA CORPORATION
 LANGUAGE OF PUBL.: English
 LANGUAGE OF FILING: English
 DOCUMENT TYPE: Patent
 PATENT INFORMATION:

	NUMBER	KIND	DATE

	WO 9742966	A1	19971120
DESIGNATED STATES:	AL AM AT AU AZ BA BB BG BR BY CA CH CN CU CZ DE DK EE ES FI GB GE GH HU IL IS JP KE KG KP KR KZ LC LK LR LS LT LU LV MD MG MK MN MW MX NO NZ PL PT RO RU SD SE SG SI SK TJ TM TR TT UA UG US UZ VN GH KE LS MW SD SZ UG AM AZ BY KG KZ MD RU TJ TM AT BE CH DE DK ES FI FR GB GR IE IT LU MC NL PT SE BF BJ CF CG CI CM GA GN ML MR NE SN TD TG		
APPLICATION INFO.:	WO 1997-US8016		19970509
PRIORITY (ORIGINAL):	US 1996-08/644287		19960510

DETD . . . culture or immunologic assays.
 Systemic
 hemodynamric signs, severe coagulopathy and intravascular thrombosis are
 notably
 absent. If properly treated, mortality is rare and neurologic sequelae,
 including
 sensineural **hearing loss**, is uncommon. The approach
 to diagnosis and
 treatment of
 meningococcal meningitis is the same as with other forms of bacterial
 meningitis.

. . .
 et al., Blood 85:3437-
 3443 (1995);
 de Winter et at., J. Inflam. 45:193-206 (1995)]. Thornton et al., FASEB
 J.,
 8(4):AI37, 1994, report that BPI **inhibited** the release of
TNF in itro
 by human
 inflammatory cells in response to LOS derived from two Neisseria species,
 N.

10 ANSWER 139 OF 142 PCTFULL COPYRIGHT 2001 MicroPatent
 ACCESSION NUMBER: 1997042966 PCTFULL
 TITLE (ENGLISH): THERAPEUTIC USES OF BPI PROTEIN PRODUCTS FOR HUMAN
 MENINGOCOCCEMIA
 TITLE (FRENCH): APPLICATIONS THERAPEUTIQUES DE PRODUITS PROTEIQUES
 BACTERICIDES/AUGMENTANT LA PERMEABILITE (BPI) DANS LE
 CAS DE
 MENINGOCOCCEMIES CHEZ L'HOMME
 INVENTOR(S): GIROIR, Brett, P.; SCANNON, Patrick, J.
 PATENT ASSIGNEE(S): XOMA CORPORATION
 LANGUAGE OF PUBL.: English
 LANGUAGE OF FILING: English
 DOCUMENT TYPE: Patent
 PATENT INFORMATION:

	NUMBER	KIND	DATE
	WO 9742966	A1	19971120
DESIGNATED STATES:	AL AM AT AU AZ BA BB BG BR BY CA CH CN CU CZ DE DK EE ES FI GB GE GH HU IL IS JP KE KG KP KR KZ LC LK LR LS LT LU LV MD MG MK MN MW MX NO NZ PL PT RO RU SD SE SG SI SK TJ TM TR TT UA UG US UZ VN GH KE LS MW SD SZ UG AM AZ BY KG KZ MD RU TJ TM AT BE CH DE DK ES FI FR GB GR IE IT LU MC NL PT SE BF BJ CF CG CI CM GA GN ML MR NE SN TD TG		
APPLICATION INFO.:	WO 1997-US8016		19970509
PRIORITY (ORIGINAL):	US 1996-08/644287		19960510

DETD . . . culture or immunologic assays.
 Systemic
 hemodynamnic signs, severe coagulopathy and intravascular thrombosis are
 notably
 absent. If properly treated, mortality is rare and neurologic sequelae,
 including
 sensineural **hearing loss**, is uncommon. The approach
 to diagnosis and
 treatment of
 meningococcal meningitis is the same as with other forms of bacterial
 meningitis.
 . . .
 et al., Blood 85:3437-
 3443 (1995);
 de Winter et at., J. Inflam. 45:193-206 (1995)]. Thornton et al., FASEB
 J.,
 8(4):AI37, 1994, report that BPI **inhibited** the release of
TNF in itro
 by human
 inflammatory cells in response to LOS derived from two Neisseria species,
 N.

L12 ANSWER 125 OF 127 PCTFULL COPYRIGHT 2001 MicroPatent
 ACCESSION NUMBER: 1993006865 PCTFULL
 TITLE (ENGLISH): TREATMENT OF OCULAR INFLAMMATION BY BLOCKAGE OF CELL
 ADHESION
 MOLECULES
 TITLE (FRENCH): TRAITEMENT DE L'INFLAMMATION OCULAIRE PAR BLOCAGE DES
 MOLECULES
 D'ADHESION CELLULAIRE
 INVENTOR(S): WHITCUP, Scott, M.; CHAN, Chi-Chao; NUSSENBLATT,
 Robert, B.
 PATENT ASSIGNEE(S): THE UNITED STATES OF AMERICA, represented by THE
 SECRETARY, DEPARTMENT
 LANGUAGE OF PUBL.: English
 DOCUMENT TYPE: Patent
 PATENT INFORMATION:

	NUMBER	KIND	DATE
	WO 9306865	A1	19930415
DESIGNATED STATES:	AU CA JP AT BE CH DE DK ES FR GB GR IE IT LU MC NL SE		
APPLICATION INFO.:	WO 1992-US8556		19921002
PRIORITY (ORIGINAL):	US 1991-770026		19911004
	US 1992-822042		19920117

DETD Case 2 A 72-year-old black woman with a 30-year
 history of bilateral recurrent panuveitis with
 exudative retinal detachments, apapilledema, **hearing**
loss, and vitiligo was diagnosed with Vogt-Koyanagi-
 Harada syndrome (Chan, C-C. et al., Am. J.

. . .
 six uveitic eyes
 showed mild staining for TNF-B (cases 1, 2, 3 and 6), and
 two uveitic eyes showed no staining with **antibody** against
TNF-B. None of the control eyes stained positively for
 either TNF-a and TNF-B.

. . .
 the regulation of
 adhesion molecule expression. The secretions of
 cytokines, particularly by the infiltrating T lymphocytes,
 probably plays an important regulatory function. Gamma-
 interferon, **interleukin-1**, and tumor necrosis factor
 cause
 the strong induction of ICAM-1, although different cells
 vary as to which cytokines will i

L12 ANSWER 125 OF 127 PCTFULL COPYRIGHT 2001 MicroPatent
 ACCESSION NUMBER: 1993006865 PCTFULL
 TITLE (ENGLISH): TREATMENT OF OCULAR INFLAMMATION BY BLOCKAGE OF CELL
 ADHESION
 MOLECULES
 TITLE (FRENCH): TRAITEMENT DE L'INFLAMMATION OCULAIRE PAR BLOCAGE DES
 MOLECULES
 D'ADHESION CELLULAIRE
 INVENTOR(S): WHITCUP, Scott, M.; CHAN, Chi-Chao; NUSSENBLATT,
 Robert, B.
 PATENT ASSIGNEE(S): THE UNITED STATES OF AMERICA, represented by THE
 SECRETARY, DEPARTMENT
 LANGUAGE OF PUBL.: English
 DOCUMENT TYPE: Patent
 PATENT INFORMATION:

	NUMBER	KIND	DATE
	WO 9306865	A1	19930415
DESIGNATED STATES:	AU CA JP AT BE CH DE DK ES FR GB GR IE IT LU MC NL SE		
APPLICATION INFO.:	WO 1992-US8556		19921002
PRIORITY (ORIGINAL):	US 1991-770026		19911004
	US 1992-822042		19920117

DETD Case 2 A 72-year-old black woman with a 30-year
 history of bilateral recurrent panuveitis with
 exudative retinal detachments, apapilledema, **hearing**
loss, and vitiligo was diagnosed with Vogt-Koyanagi-
 Harada syndrome (Chan, C-C. et al., Am. J.

. . .
 six uveitic eyes
 showed mild staining for TNF-B (cases 1, 2, 3 and 6), and
 two uveitic eyes showed no staining with **antibody** against
TNF-B. None of the control eyes stained positively for
 either TNF-a and TNF-B.

. . .
 the regulation of
 adhesion molecule expression. The secretions of
 cytokines, particularly by the infiltrating T lymphocytes,
 probably plays an important regulatory function. Gamma-
 interferon, **interleukin-1**, and tumor necrosis factor
 cause
 the strong induction of ICAM-1, although different cells
 vary as to which cytokines will i

L12 ANSWER 122 OF 127 PCTFULL COPYRIGHT 2001 MicroPatent
 ACCESSION NUMBER: 1999053927 PCTFULL
 TITLE (ENGLISH): METHODS FOR TREATING AND PREVENTING INSULIN RESISTANCE
 AND
 RELATED DISORDERS
 TITLE (FRENCH): PROCEDES POUR TRAITER ET PREVENIR LA RESISTANCE A
 L'INSULINE ET
 LES TROUBLES QUI Y SONT LIES
 INVENTOR(S): GREENBERG, Andrew, S.
 PATENT ASSIGNEE(S): TRUSTEES OF TUFTS COLLEGE
 LANGUAGE OF PUBL.: English
 LANGUAGE OF FILING: English
 DOCUMENT TYPE: Patent
 PATENT INFORMATION:

	NUMBER	KIND	DATE
	WO 9953927	A1	19991028
DESIGNATED STATES:	JP US AT BE CH CY DE DK ES FI FR GB GR IE IT LU MC NL		
	PT SE		
APPLICATION INFO.:	WO 1999-US8364		19990416
PRIORITY (ORIGINAL):	US 1998-60/082152		19980417
	US 1998-		19980423

DETD . . . years while attacking the blood
 vessels and nerves,
 Diabetics, as a group, are far more often afflicted with blindness,
 heart disease, stroke, kidney
 disease, **hearing loss**, gangrene and impotence. One
 third of all visits
 to physicians are
 occasioned by this disease and its complications, and diabetes and its
 complications. . .

. . .
 activity of JNK and also, but to a
 lesser extent, the activity
 of ERK1" kinase. Furthermore, a prostaglandin derivative was also shown
 to **inhibit TNF-**
 u induced lipolysis and to **inhibit** ERK1/2 and YNK-1 activation.
 Furthermore, as shown
 herein, an inhibitor of a ERK/MA-P kinase decreases TNF-a induced
 lipolysis. In addition,
 contrary to JNK and. . .) 8 pathway
 stimulates TNF-a
 induced lipolysis. Thus, the Examples described herein indicate that
 inhibition of the JNK
 pathway and the ERK/MAP kinase pathway **inhibit TNF-(y**
 induced lipolysis,
 whereas
inhibition of the p3) 8 pathway increases TNF-a induced
 lipolysis.

3. Compounds which **inhibit TNF-a** induced lipolysis
 The invention provides methods for inhibiting, or blocking lipolysis of
 adipocytes. In
 In
 a preferred embodiment, lipolysis is TNF-a induced lipolysis. In. . .

. . .
 kinases
 (SAPKs), are
 members of the mitogen-activated protein (MAP) kinase group which are
 activated in
 response to cytokines, such as TNF, e.g., TNF-(x and **IL-**
1, and exposure
 to environmental
 ZD
 stress, including ultraviolet light, heat shock, and osmotic stress (U.S.
 Pat. No. 5605808; B.

Other preferred compounds of the invention include compounds which inhibit TNF- α .

Such compounds include those which inhibit the production of TNF-a, as well as those which specifically inhibit its activity, e.g., by interfering with its interaction with a receptor. For example, an inhibitor can be an antibody binding specifically to, and inhibiting the activity of TNF-a. Yet other **inhibitory** compounds include **inhibitors** of a TNF-a receptor or of a molecule interacting therewith, such as the NLADD protein (Schievella et al. (1997) J Biol, Chem. 272:12069). Thus, within the scope of the invention are antisense and triplex molecules; which specifically **inhibit** the expression of **TNF-a**, a TNF-u receptor, or molecule interacting with TNF-u, or such as MADD.

thereof As described in the Examples, NaSal has in fact been shown herein to inhibit JNK activation by TNF-u. in adipocytes and to **inhibit TNF**-cz induced lipolysis in adlpoc tes. NaSal has also been described as being capable of **inhibiting TNF**-Induced activation of JNK and p42/p44 MAP kinases in FS4 fibroblasts (Schwenger et al. (1997) P.N.A.S. tWA 94:2869) and Schwenger et al. (1996). . .

discovery that the TNK and the EPK/MAP kinase pathways are involved in TNF- α induced lipolysis, and that inhibition of these kinase pathways decreases or **inhibits** TNF- α induced lipolysis. It has also been shown that PPAR- γ agonists inhibit TNF- α induced lipolysis. Thus, the invention pertains to any disease or disorder which. . . .

In a particular embodiment, insulin resistance or other disease or condition associated with an abnormal FFA level is reduced, eliminated or **inhibited**, by blocking **TNF**- α action, e.g., by neutralizing TNF- α in the serum (i.e., circulating TNF- α) or in the adipose tissue.

methods of the invention comprise determining the activity of a PPAR- γ receptor, In fact, since binding of a ligand to a PPAR- γ receptor **inhibits** **TNF**- α induced lipolysis, a defective PPAR- γ may result in increased TNF- α induced lipolysis.

of detection is by performing ELISA assays. An ELISA assay for the detection of human T.NF-rt, as well as the production of **antibodies** against human **TNF-** U., is described, e.g., in U.S. Patent No. 5,716,972 by Adams et al. Antibodies for performing

these assays are also commercially available, such as. . .

shown in Figure 1. These indicate that lipolysis in the adipocytes induced by 2 ng/ml (Panel A) or 20 ng/ml (Panel B) **TNF- α** is significantly **inhibited** by the addition of 2 or 20 mM NaSal. In fact, NaSal inhibited glycerol production by 3T3 LI adipocytes induced by. . .

Example 2: NaSal **inhibits** **TNF- α** induced JNK-1 activation in adipocytes
This example shows that NaSal inhibits JNK-1 activation by **TNF- α** in adipocytes.

Thus, since NaSal **inhibits** **TNF- α** induced lipolysis and NaSal significantly reduces JNK-1 activation, JNK-1 activation is likely to play a significant role in blocking lipolysis induced by **TNF- α** .

Example 3. NaSal partially **inhibits** **TNF- α** induced NMP kinase activation
This example shows that, whereas JNK-1 activation by **TNF- α** is significantly **inhibited** by NaSal, activation of NALP kinases by **TNF- α** is not significantly inhibited by NaSal.

Example 4- NMP kinase **inhibitor** PD98059 decreases **TNF- α** induced lipolysis
This example demonstrates that inhibition of ERK1/2 by PD98059 decreases **TNF- α** induced lipolysis.

Example 5: **Inhibition** of p38 kinase stimulates **TNF- α** induced lipolysis
This example shows; that, contrarily to JNK and ERK 1/2 signal pathways, inhibition of the p38 signal pathway stimulates **TNF- α** induced. . .

ERK 1/2 kinase activity, BRL does not significantly affect the activity of JNK-1 and ERK 1/2 kinases. Thus, BRL does not **inhibit** **TNF- α** induced lipolysis by decreasing the activity of JNK-1 and ERK.

Example 7: PGJ2 **inhibits** **TNF- α** induced activation of JNK-1 and ERK1/2
This example shows that **TNF- α** induced activation of JNK-1 and ERK1/2 can be inhibited by PGJ2.

p38. Total protein is shown in Figure 7 panel B. Furthermore, Figure 7 panels C and D shows that PGJ2 significantly **inhibits** the **TNF- α** induced increase in ERK 1/2 and that of JNK-1 kinase activity. Thus., PGJ2, which is an agonist of the PPAR- γ receptor may function in a similar manner to NaSal to **inhibit** **TNF- α** induced lipolysis, i.e., by modulating TVAP kinases, e.g., ERK1/2 and JNK-1.

Example 8: **Inhibition** of JNK-1 decreases **TNF-u**,
induced lipolysis

This example demonstrates an experiment that a person of skill in the
art can effectuate
without undue experimentation to show that. . .

adenoviral vectors. Furthermore, as shown in Figure I OA,
incubation of the cells
with all adenoviral vector expressing perilipin A or B, essentially
inhibited completely lipolysis
induced by **TNF-a**, whereas the adenoviral vector alone had no
effect on
TNF-u. induced
lipolysis. As shown in Figure I OB, **TNF-r4** induced lipolysis
was still

inhibited during the last ')
hours of the 24 hour treatment. On the contrary, perilipin A or B
overexpression did not
significantly block. . .

L12 ANSWER 122 OF 127 PCTFULL COPYRIGHT 2001 MicroPatent
 ACCESSION NUMBER: 1999053927 PCTFULL
 TITLE (ENGLISH): METHODS FOR TREATING AND PREVENTING INSULIN RESISTANCE
 AND
 RELATED DISORDERS
 TITLE (FRENCH): PROCEDES POUR TRAITER ET PREVENIR LA RESISTANCE A
 L'INSULINE ET
 LES TROUBLES QUI Y SONT LIES
 INVENTOR(S): GREENBERG, Andrew, S.
 PATENT ASSIGNEE(S): TRUSTEES OF TUFTS COLLEGE
 LANGUAGE OF PUBL.: English
 LANGUAGE OF FILING: English
 DOCUMENT TYPE: Patent
 PATENT INFORMATION:

	NUMBER	KIND	DATE
	WO 9953927	A1	19991028
DESIGNATED STATES:	JP US AT BE CH CY DE DK ES FI FR GB GR IE IT LU MC NL		
	PT SE		
APPLICATION INFO.:	WO 1999-US8364		19990416
PRIORITY (ORIGINAL):	US 1998-60/082152		19980417
	US 1998-		19980423

DETD . . . years while attacking the blood
 vessels and nerves,
 Diabetics, as a group, are far more often afflicted with blindness,
 heart disease, stroke, kidney
 disease, **hearing loss**, gangrene and impotence. One
 third of all visits
 to physicians are
 occasioned by this disease and its complications, and diabetes and its
 complications. . .

. . .
 activity of JNK and also, but to a
 lesser extent, the activity
 of ERK1" kinase. Furthermore, a prostaglandin derivative was also shown
 to **inhibit TNF-**
 u induced lipolysis and to **inhibit** ERK1/2 and YNK-1 activation.
 Furthermore, as shown
 herein, an inhibitor of a ERK/MA-P kinase decreases TNF-a induced
 lipolysis. In addition,
 contrary to JNK and. . .) 8 pathway
 stimulates TNF-a
 induced lipolysis. Thus, the Examples described herein indicate that
 inhibition of the JNK
 pathway and the ERK/MAP kinase pathway **inhibit TNF-**(y
 induced lipolysis,
 whereas
inhibition of the p3) 8 pathway increases TNF-a induced
 lipolysis.

3. Compounds which **inhibit TNF-**a induced lipolysis
 The invention provides methods for inhibitino, or blocking lipolysis of
 adipocytes. In
 In
 a preferred embodiment, lipolysis is TNF-a induced lipolysis. In. . .

. . .
 kinases
 (SAPKs), are
 members of the mitogen-activated protein (MAP) kinase group which are
 activated in
 response to cytokines, such as TNF, e.g., TNF-(x and **IL-**
1, and exposure
 to environmental
 ZD
 stress, including ultraviolet light, heat shock, and osmotic stress (U.S.
 Pat. No. 5605808; B.

Other compounds of the invention
Other preferred compounds of the invention include compounds which
inhibit TNF-a.

Such compounds include those which inhibit the production of TNF-a, as well as those which specifically inhibit its activity, e.g., by interfering. . . its interaction With a receptor. For example, an inhibitor can be an antibody binding specifically to, and inhibiting the activity of TNF-a. Yet other **inhibitory** compounds include **inhibitors** of a TNF-a receptor or of a molecule interacting therewith, such as the NLADD protein (Schievella et al. (1997) J Biol, Chem. 272:12069). Thus, within the scope of the invention are antisense and triplex molecules; which specifically **inhibit** the expression of TNF-a, a TNF-u. receptor, or molecule interactina with TNF-u, or' uch as MADD.

thereof As described in the Examples,
NaSal has in fact been shown herein to inhibit JNK activation by TNF-u. in adipocytes and to **inhibit TNF-cz** induced lipolysis in adlpoc tes. NaSal has also been described as being capable of **inhibiting** TNF-Induced activation of JNK and p42/p44 MAP kinases in FS4 fibroblasts (Schwenger et al. (1997) P.N.A.S. tWA 94:2869) and Schwenger et al. (1996). . .

discovery that the.TNK and the EPK/MAP kinase pathways are involved in TNF-a induced lipolysis, and that iri-hibition of these kinase pathways decreases or **inhibits** TNF-a induced lipolysis. It has also been shown that PPAR-y agonists inhibit TNIF-a induced lipolysis. Thus, the invention pertains to any disease or dlsorder which. . .

In a particular embodiment, insulin resistance or other disease or condition associated with an abnormal FFA level is reduced, eliminated or **inhibited**, by blocking TNF-a action, e-, (,- by neutralizing TNF-u, in the serum (Le., circulating T'NT,-ct) or in the adipose tissue.

methods of the invention comprise determining the activity of a PPAR-y receptor, In fact, since binding of a ligand to a PPAR-y receptor **inhibits** TNF-a induced lipolysis, a defective PPAR-y may result in increased TNF-a induced lipolysis.

of detection is by performing ELISA assays. An ELISA assay for the detection of human T.NF-rt, as well as the production of **antibodies** against human TNF-U., is described, e.g., in U.S. Patent No. 5,716,972 by Adams eL a]. Antibodies for performing

these assays are also commercially available, such as. . .

showed in Figure 1. These indicate that lipolysis in the adipocytes induced by 2 ng/ml (Panel A) or 20 ng/ml (Panel B) **TNF- α** is significantly **inhibited** by the addition of 2 or 20 mM NaSal. In fact, NaSal inhibited glycerol production by 3T3 LI adipocytes induced by. . .

Example 2: NaSal **inhibits** **TNF- α** induced JNK-1 activation in adipocytes
This example shows that NaSal inhibits JNK-1 activation by **TNF- α** in adipocytes.

Thus, since NaSal **inhibits** **TNF- α** induced lipolysis and NaSal significantly reduces JNK-1 activation, JNK-1 activation is likely to play a significant role in blocking lipolysis induced by **TNF- α** .

Example 3. NaSal partially **inhibits** **TNF- α** induced NMP kinase activation
This example shows that, whereas JNK-1 activation by **TNF- α** is significantly **inhibited** by NaSal, activation of NALP kinases by **TNF- α** is not significantly inhibited by NaSal.

Example 4- NMP kinase **inhibitor** PD98059 decreases **TNF- α** induced lipolysis
This example demonstrates that inhibition of ERK1/2 by PD98059 decreases **TNF- α** induced lipolysis.

Example 5: **Inhibition** of p38 kinase stimulates **TNF- α** induced lipolysis
This example shows; that, contrarily to JNK and ERK 1/2 signal pathways, inhibition of the p38 signal pathway stimulates **TNF- α** induced. . .

ERK 1/2 kinase activity, BRL does not significantly affect the activity of JNK-1 and ERK 1/2 kinases. Thus, BRL does not **inhibit** **TNF- α** induced lipolysis by decreasing the activity of JNK-1 and ERK.

Example 7: PGJ2 **inhibits** **TNF- α** induced activation of JNK-1 and ERK1/2
This example shows that **TNF- α** induced activation of JNK-1 and ERK1/2 can be inhibited by PGJ2.

p38. Total protein is shown in Figure 7 panel B. Furthermore, Figure 7 panels C and D shows that PGJ2 significantly **inhibits** the **TNF- α** induced increase in ERK 1/2 and that of JNK-1 kinase activity. Thus., PGJ2, which is an agonist of the PPAR- γ receptor may function in a similar manner to NaSal to **inhibit** **TNF- α** induced lipolysis, i.e., by modulating MAP kinases, e.g., ERK1/2 and JNK-1.

Example 8: **Inhibition** of JNK-1 decreases **TNF-u**,
induced lipolysis

This example demonstrates an experiment that a person of skill in the
art can effectuate
without undue experimentation to show that. . .

adenoviral vectors. Furthermore, as shown in Figure I OA,
incubation of the cells
with all adenoviral vector expressing perilipin A or B, essentially
inhibited completely lipolysis
induced by **TNF-a**, whereas the adenoviral vector alone had no
effect on
TNF-u. induced
lipolysis. As shown in Figure I OB, **TNF-r4** induced lipolysis
was still

inhibited during the last ')
hours of the 24 hour treatment. On the contrary, perilipin A or B
overexpression did not
significantly block. . .

WEST[Help](#)[Logout](#)[Interrupt](#)[Main Menu](#)[Search Form](#)[Posting Counts](#)[Show S Numbers](#)[Edit S Numbers](#)[Preferences](#)**Search Results -**

Term	Documents
MENINGITIS.DWPI,EPAB,JPAB,USPT,PGPB.	2980
MENINGITI.DWPI,EPAB,JPAB,USPT,PGPB.	5
HEARING.DWPI,EPAB,JPAB,USPT,PGPB.	21362
HEARINGS.DWPI,EPAB,JPAB,USPT,PGPB.	233
LOSS.DWPI,EPAB,JPAB,USPT,PGPB.	655519
LOSSES.DWPI,EPAB,JPAB,USPT,PGPB.	164374
(MENINGITIS SAME (HEARING ADJ LOSS)).USPT,PGPB,JPAB,EPAB,DWPI.	16

US Patents Full-Text Database
US Pre-Grant Publication Full-Text Database
JPO Abstracts Database
EPO Abstracts Database
Derwent World Patents Index

Database:

IBM Technical Disclosure Bulletins

meningitis same (hearing adj loss)

Refine Search:**Clear****Search History****Today's Date: 9/20/2001**

WEST[Help](#)[Logout](#)[Interrupt](#)[Main Menu](#)[Search Form](#)[Posting Counts](#)[Show S Numbers](#)[Edit S Numbers](#)[Preferences](#)**Search Results -**

Term	Documents
MENINGITIS.DWPI,EPAB,JPAB,USPT,PGPB.	2980
MENINGITI.DWPI,EPAB,JPAB,USPT,PGPB.	5
HEARING.DWPI,EPAB,JPAB,USPT,PGPB.	21362
HEARINGS.DWPI,EPAB,JPAB,USPT,PGPB.	233
LOSS.DWPI,EPAB,JPAB,USPT,PGPB.	655519
LOSSES.DWPI,EPAB,JPAB,USPT,PGPB.	164374
(MENINGITIS SAME (HEARING ADJ LOSS)).USPT,PGPB,JPAB,EPAB,DWPI.	16

US Patents Full-Text Database
US Pre-Grant Publication Full-Text Database
JPO Abstracts Database
EPO Abstracts Database
Derwent World Patents Index

Database:

IBM Technical Disclosure Bulletins

meningitis same (hearing adj loss)

Refine Search:[Clear](#)**Search History****Today's Date: 9/20/2001**

<u>DB Name</u>	<u>Query</u>	<u>Hit Count</u>	<u>Set Name</u>
USPT,PGPB,JPAB,EPAB,DWPI	meningitis same (hearing adj loss)	16	<u>L6</u>
USPT,PGPB,JPAB,EPAB,DWPI	meningitis adj5 (hearing adj loss)	1	<u>L5</u>
USPT,PGPB,JPAB,EPAB,DWPI	meningitis adj5(hearing adj loss)	1	<u>L4</u>
USPT,PGPB,JPAB,EPAB,DWPI	(TNF or tumor adj necro\$3 adj factor or tumour adj necro\$4 adj factor) same (il-1 or interleukin-1) and (hearing adj loss or cochlear adj degenrat\$)	13	<u>L3</u>
USPT,PGPB,JPAB,EPAB,DWPI	(TNF or tumor adj necro\$3 adj factor or tumour adj necro\$4 adj factor) same meningitis and (hearing adj loss or cochlear adj degenrat\$)	4	<u>L2</u>
USPT,PGPB,JPAB,EPAB,DWPI	(TNF or tumor adj necro\$3 adj factor or tumour adj necro\$4 adj factor) same (hearing adj loss or cochlear adj degenrat\$)	4	<u>L1</u>

<u>DB Name</u>	<u>Query</u>	<u>Hit Count</u>	<u>Set Name</u>
USPT,PGPB,JPAB,EPAB,DWPI	meningitis same (hearing adj loss)	16	<u>L6</u>
USPT,PGPB,JPAB,EPAB,DWPI	meningitis adj5 (hearing adj loss)	1	<u>L5</u>
USPT,PGPB,JPAB,EPAB,DWPI	meningitis adj5(hearing adj loss)	1	<u>L4</u>
USPT,PGPB,JPAB,EPAB,DWPI	(TNF or tumor adj necro\$3 adj factor or tumour adj necro\$4 adj factor) same (il-1 or interleukin-1) and (hearing adj loss or cochlear adj degenrat\$)	13	<u>L3</u>
USPT,PGPB,JPAB,EPAB,DWPI	(TNF or tumor adj necro\$3 adj factor or tumour adj necro\$4 adj factor) same meningitis and (hearing adj loss or cochlear adj degenrat\$)	4	<u>L2</u>
USPT,PGPB,JPAB,EPAB,DWPI	(TNF or tumor adj necro\$3 adj factor or tumour adj necro\$4 adj factor) same (hearing adj loss or cochlear adj degenrat\$)	4	<u>L1</u>